

Giant cockroach における新規糖脂質の構造研究

Michael K. STOSKOPF^{*1}, Yasuo KISHIMOTO^{*2}, Takashi TANAKA^{*3},
岡村信幸, Lou-sing Kan^{*4}, Robert COTTER^{*4}, Catherine FENSELAU^{*5}

Journal of Biological Chemistry, 264 (9), 4964-4971 (1989)

Characterization of Novel Glycolipids from the Giant Cockroach (*Blaberus colosseus*)

Michael K. STOSKOPF^{*1}, Yasuo KISHIMOTO^{*2}, Takashi TANAKA^{*3},
Nobuyuki OKAMURA, Lou-sing Kan^{*4}, Robert COTTER^{*4},
Catherine FENSELAU^{*5}

ABSTRACT A novel class of glycolipids, assigned the trivial name blaberosides, was isolated from whole head tissues of the giant cockroach (*Blaberus colosseus*). The class consists of two closely related families, blaberoside I and blaberoside II, each containing species differing by 26 atomic mass units. The structure of these gentiobiose-based glycosylglycerolipids was elucidated by chromatographic behavior, nuclear magnetic resonance spectroscopy, mass spectrometry, and analysis of chemical degradation products and derivatives. Species in the blaberoside I family have been identified as 2-O-[6'-O-(6''-O-3-hydroxy-11-eicosenoyl- β -D-glucopyranosyl)- β -D-glucopyranosyl]-3-(hexadecyloxy)-1-(3-hydroxy-11-eicosenoyl)-1, 2-propanediol (blaberoside Ia) and 2-O-[6'-O-(6''-O-3-hydroxy-11-eicosenoyl- β -D-glucopyranosyl)- β -D-glucopyranosyl]-3-(6-octadecyloxy)-1-(3-hydroxy-11-eicosenoyl)-1, 2-propanediol (blaberoside Ib). Two smaller homologs of the blaberoside II family were discerned to be 2-O-[6'-O-(6''-O-3-hydroxy-11-eicosenoyl- β -D-glucopyranosyl)- β -D-glucopyranosyl]-3-(hexadecyloxy)-1, 2-propanediol (blaberoside IIa), and 2-O-[6'-O-(6''-O-3-hydroxy-11-eicosenoyl- β -D-glucopyranosyl)- β -D-glucopyranosyl]-3-(4-octadecyloxy)-1, 2-propanediol (blaberoside IIb). These compounds are unique because they are animal origin glycosylglycerolipids with a highly flexible gentiobiose backbone, and a β -linkage of the carbohydrate to the glycerol ether at the 2 position rather than the usual 1 position.

抄録 giant cockroach (*Blaberus colosseus*) の頭部組織から新しい種類の糖脂質である blaberoside I と blaberoside II を単離した。構造研究の結果, これらはゲンチオビオースからなる糖脂質で, いずれも分子量が26異なった2つの化合物から構成され, blaberoside I は blaberoside Ia と Ib, blaberoside II は blaberoside IIa と IIb であると決定した。これらの化合物は, 柔軟性に富んだ糖の結合様式を有するゲンチオビオースを基本骨格としていることと, 通常と異なりグリセロールの2位に糖が β 結合したグリセロ糖脂質であるという点で非常にめずらしい。

*1 : School of Veterinary Medicine, North Carolina State University

ノースカロライナ州立大学獣医学部

*2 : School of Medicine, University of California, San Diego

カリフォルニア大学医学部サンディエゴ校

*3 : 九州大学薬学部

*4 : School of Medicine, Johns Hopkins University ジョーンズ ホプキンス大学医学部

*5 : University of Maryland メリーランド大学